

## **INFORMED CONSENT - BLEHPAROPLASTY SURGERY**

### **INSTRUCTIONS**

This is an informed-consent document which has been prepared to help inform you about blepharoplasty (eyelid) surgery, its risks, as well as alternative treatment(s).

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon and agreed upon by you.

Blepharoplasty is a surgical procedure to remove excess skin and muscle from both the upper and lower eyelids. Underlying fatty tissue that produces bagginess can be selectively removed or repositioned. Blepharoplasty can improve drooping skin and bagginess. It can help improve vision in older patients who have hooding of their upper eyelids. Although it can add an upper eyelid crease to the Asian eyelid, it will not erase evidence of one's racial or ethnic heritage. Blepharoplasty will not remove "crow's feet" or other wrinkles, eliminate dark circles under the eyes, or lift sagging eyebrows.

Blepharoplasty surgery is customized for every patient, depending on his or her particular needs. It can be performed alone involving upper, lower or both eyelid regions, or in conjunction with other surgical procedures of the eye, face, brow, or nose. In some patients, who have looseness between the lower eyelid and the eyeball, consideration for tightening of the lower eyelid (canthoplasty/canthopexy) at the time of blepharoplasty may be recommended. Eyelid surgery cannot stop the process of aging. It can however, diminish the look of loose skin and bagginess in the eyelid region.

### **ALTERNATIVE TREATMENTS**

Alternative forms of management include not treating the skin laxness and bagginess in the eyelids by surgery. Improvement of skin laxness, fatty deposits and skin wrinkles may be accomplished by other treatments or surgery such as a brow lift when indicated. Other forms of eyelid surgery may be needed should you have disorders affecting the function of the eyelid such as drooping eyelids from muscle problems (eyelid ptosis) or looseness between the eyelid and eyeball (ectropion). Minor skin wrinkling may be improved through chemical skin-peels, laser resurfacing, or other skin treatments. Risks and potential complications are associated with alternative surgical forms of treatment.

### **RISKS OF BLEPHAROPLASTY SURGERY**

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations.

An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your plastic surgeon to make sure you understand all possible consequences of

**Bleeding-** It is possible, though unusual, to experience a bleeding episode during or after surgery. Bleeding may occur under the skin or internally around the eyeball. Intraoperative blood transfusions may be required. Should post-operative bleeding occur, it may require emergency treatment, surgery, or blood transfusion. Do not take any aspirin or anti-inflammatory medications for ten days before or after surgery, as this may increase the risk of bleeding. Nonprescription "herbs" and dietary supplements can increase the risk of surgical bleeding. Hypertension (high blood pressure) that is not under good medical control may cause bleeding during or after surgery. Accumulations of blood under the eyelids may delay healing and cause scarring. Hematoma can occur at any time following injury. If blood transfusions are needed to treat blood loss, there is a risk of blood-related infections such as hepatitis and HIV (AIDS). Heparin medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

## INFORMED CONSENT - BLEHPAROPLASTY SURGERY

**Blindness-** Blindness is extremely rare after blepharoplasty. However, it can be caused by internal bleeding around the eye during or after surgery. The occurrence of this is not predictable.

**Infection-** Infection is unusual after surgery. Should an infection occur, additional treatment including antibiotics, hospitalization, or additional surgery may be necessary.

**Scarring-** All surgery leaves scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the eyelid and deeper tissues. Scars may be unattractive and of different color than the surrounding skin tone. Scar appearance may also vary within the same scar. Scars may be asymmetrical (appear different on the right and left side of the body). There is the possibility of visible marks in the eyelid or small skin cysts from sutures. In some cases

**Damage to Deeper Structures-** There is the potential for injury to deeper structures including, nerves, blood vessels, eye muscles, and lungs (pneumothorax) during any surgical procedure. The potential for this to occur varies according to the type of blepharoplasty procedure performed. Injury to deeper

**Dry Eye Problems-** Permanent disorders involving decreased tear production can occur after blepharoplasty. The occurrence of this is rare and not entirely predictable. Individuals who normally have dry eyes may be advised to use special caution in considering blepharoplasty surgery.

**Asymmetry-** The human face is normally asymmetrical. There can be a variation from one side to the other in the results obtained from blepharoplasty surgery. Additional surgery may be necessary to attempt

**Pain-** You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after blepharoplasty surgery. Chronic pain may occur very infrequently from nerves becoming

**Ectropion-** Displacement of the lower eyelid away from the eyeball is a rare complication. Further surgery

**Corneal Exposure Problems-** Some patients experience difficulties closing their eyelids after surgery and problems may occur in the cornea due to dryness. Should this rare complication occur, additional treatments or surgery and treatment may be necessary.

**Allergic Reactions-** In rare cases, local allergies to tape, suture materials and glues, blood products, topical preparations or injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur to drugs used during surgery and prescription medications. Allergic reactions

**Eyelash Hair Loss-** Hair loss may occur in the lower eyelash area where the skin was elevated during surgery. The occurrence of this is not predictable. Hair loss may be temporary or permanent.

**Delayed Healing-** Wound disruption or delayed wound healing is possible. Some areas of the skin may not heal normally and may take a long time to heal. Areas of skin may die. This may require frequent dressing changes or further surgery to remove the non-healed tissue. Smokers have a greater risk of skin loss and wound healing complications.

**Change in Skin Sensation-** It is common to experience diminished (or loss) of skin sensation in areas that have had surgery. Diminished (or complete loss of skin sensation) may not totally resolve after a blepharoplasty.

## **INFORMED CONSENT - BLEHPAROPLASTY SURGERY**

**Skin Contour Irregularities-** Contour irregularities and depressions may occur after blepharoplasty. Visible and palpable wrinkling of skin can occur. Residual skin irregularities at the ends of the incisions or “dog ears” are always a possibility and may require additional surgery. This may improve with time, or it can be surgically corrected.

**Skin Discoloration / Swelling-** Some bruising and swelling normally occurs following blepharoplasty. The skin in or near the surgical site can appear either lighter or darker than surrounding skin. Although uncommon, swelling and skin discoloration may persist for long periods of time and, in rare situations,

**Skin Sensitivity-** Itching, tenderness, or exaggerated responses to hot or cold temperatures may occur after surgery. Usually this resolves during healing, but in rare situations it may be chronic.

**Sutures-** Most surgical techniques use deep sutures. You may notice these sutures after your surgery. Sutures may spontaneously poke through the skin, become visible or produce irritation that requires

**Surgical Anesthesia-** Both local and general anesthesia involve risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

**Shock-** In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment would be necessary.

**Seroma-** Fluid accumulations infrequently occur in between the skin and the underlying tissues. Should this problem occur, it may require additional procedures for drainage of fluid.

**Unsatisfactory Result-**Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. You may be disappointed with the results of eyelid surgery. This would include risks such as asymmetry, unsatisfactory surgical scar location, unacceptable visible deformities, loss of function, poor healing, wound disruption, and loss of sensation.

It may be necessary to perform additional surgery to improve your results. Additional surgical procedures such as a brow lift may be needed to correct eyebrow sagging which contributes to upper eyelid

**Cardiac and Pulmonary Complications-** Surgery, especially longer procedures, may be associated with the formation of, or increase in, blood clots in the venous system. Pulmonary complications may occur secondarily to both blood clots (pulmonary emboli), fat deposits (fat emboli) or partial collapse of the lungs after general anesthesia. Pulmonary and fat emboli can be life-threatening or fatal in some circumstances. Air travel, inactivity and other conditions may increase the incidence of blood clots traveling to the lungs causing a major blood clot that may result in death. It is important to discuss with your physician any past history of blood clots or swollen legs that may contribute to this condition.

Cardiac complications are a risk with any surgery and anesthesia, even in patients without symptoms. If you experience shortness of breath, chest pains, or unusual heart beats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional

### **ADDITIONAL ADVISORIES**

**Long-Term Results-** Subsequent alterations in eyelid appearance may occur as the result of aging, weight loss or gain, sun exposure, pregnancy, menopause, or other circumstances not related to blepharoplasty surgery. Blepharoplasty surgery does not arrest the aging process or produce permanent tightening of the eyelid region. Future surgery or other treatments may be necessary to maintain the results of a blepharoplasty.

## INFORMED CONSENT - BLEHPAROPLASTY SURGERY

**Skin Disorders / Skin Cancer-** A blepharoplasty is a surgical procedure to tighten the loose skin and deeper structures of the eyelid. Skin disorders and skin cancer may occur independently of eyelid surgery.

**Female Patient Information-** It is important to inform your plastic surgeon if you use birth control pills, estrogen replacement, or if you believe you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

**Intimate Relations After Surgery-** Surgery involves coagulating of blood vessels and increased activity of any kind may open these vessels leading to a bleed, or hematoma. Activity that increases your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery and control bleeding. It is wise to refrain from sexual activity until your physician states it is safe.

**Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray)-** Patients who are currently smoking, use tobacco products, or nicotine products (patch, gum, or nasal spray) are at a greater risk for significant surgical complications of skin dying, delayed healing, and additional scarring. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of this type of complication. Please indicate your current status regarding these items below: I am a non-smoker and do not use nicotine products. I understand the risk of second-hand

\_\_\_ ^^ I am a smoker or use tobacco / nicotine products. I understand the risk of surgical

It is important to refrain from smoking at least 6 weeks before surgery and until your physician states it is

**Mental Health Disorders and Elective Surgery-** It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less than satisfactory results are sometimes unavoidable, may require additional surgery and often are stressful. Please openly discuss with your surgeon, prior to surgery, any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately

**Medications-** There are many adverse reactions that occur as the result of taking over-the-counter, herbal, and/or prescription medications. Be sure to check with your physician about any drug interactions that may exist with medications which you are already taking. If you have an adverse reaction, stop the drugs immediately and call your plastic surgeon for further instructions. If the reaction is severe, go immediately to the nearest emergency room. When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, do not operate complex equipment, do not make any important decisions, and do not drink any alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

### **ADDITIONAL SURGERY NECESSARY**

There are many variable conditions in addition to risk and potential surgical complications that may influence the long-term result of eyelid surgery. Secondary surgery may be necessary to obtain optimal results. Even though risks and complications occur infrequently, the risks cited are the ones that are particularly associated with blepharoplasty surgery. Other complications and risks can occur but are even more uncommon. Should complications occur, additional surgery or other treatments may be necessary. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. In some situations, it may not be possible to achieve optimal results with a single surgical procedure.

## **INFORMED CONSENT - BLEHPAROPLASTY SURGERY**

### **PATIENT COMPLIANCE**

Follow all physician instructions carefully; this is essential for the success of your outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activity needs to be restricted. Protective dressings and drains should not be removed unless instructed by your plastic surgeon. Successful post-operative function depends on both surgery and subsequent care. Physical activity that increases your pulse or heart rate may cause bruising, swelling, fluid accumulation and the need for return to surgery. It is wise to refrain from intimate physical activities after surgery until your physician states it is safe. It is important that you participate in follow-up care, return for aftercare, and promote your recovery after surgery.

If hooding of the upper eyelids interfere with your vision, your health insurance company may cover blepharoplasty surgery for the upper-eyelids only. Most health insurance companies exclude coverage for cosmetic surgical operations such as the lower-eyelid blepharoplasty or any complications that might occur from surgery. Most insurance plans exclude coverage for secondary or revisionary surgery. Please carefully review your health insurance subscriber information pamphlet.

### **FINANCIAL RESPONSIBILITIES**

The cost of surgery involves several charges for the services provided. The total includes fees charged by your surgeon, the cost of surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day-surgery charges involved with revision surgery will also be your responsibility. In signing the consent for this surgery/procedure, you acknowledge that you have been informed about its risk and consequences and accept responsibility for the clinical decisions that were made along with the financial costs of all future treatments.

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s), including no surgery. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed-consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information which is based on all the facts in your particular case and the current state of medical

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

**INFORMED CONSENT - BLEHPAROPLASTY SURGERY**

**CONSENT FOR SURGERY / PROCEDURE or TREATMENT**

1. I hereby authorize Dr. \_\_\_\_\_ and such assistants as may be selected to perform the following procedure or treatment:

EYELID SURGERY (Blepharoplasty)

I have received the following information sheet:

INFORMED CONSENT - BLEPHAROPLASTY SURGERY

2. I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants or designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure is begun.

3.1 consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and the possibility of complications, injury, and sometimes death.

4.1 acknowledge that no guarantee or representation has been given by anyone as to the results that may be obtained.

5.1 consent to be photographed or televised before, during, and after the operation(s) or procedure(s) to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.

6. For purposes of advancing medical education, I consent to the admittance of observers to the operating room.

7.1 consent to the disposal of any tissue, medical devices or body parts which may be removed.

8.1 consent to the utilization of blood products should they be deemed necessary by my surgeon and/or his/her appointees, and I am aware that there are potential significant risks to my health with their utilization.

9.1 authorize the release of my Social Security number to appropriate agencies for legal reporting and medical-device registration, if applicable.

10.1 understand that the surgeons' fees are separate from the anesthesia and hospital charges, and the fees are agreeable to me. If a secondary procedure is necessary, further expenditure will be required.

11.1 realize that not having the operation is an option.

12. IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:

a. THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN

b. THERE MAY BE ALTERNATIVE PROCEDURES OR METHODS OF TREATMENT

c. THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

I CONSENT TO THE TREATMENT OR PROCEDURE AND THE ABOVE LISTED ITEMS (1-

12). I AM SATISFIED WITH THE EXPLANATION.

Patient or Person Authorized to Sign for Patient

Date \_\_\_\_\_ Witness